

Tilburg University

On the estimated variances of regression coefficients in misspecified error components models

Deschamps, P.J.

Publication date:
1992

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Tilburg University Research Portal](#)

Citation for published version (APA):
Deschamps, P. J. (1992). *On the estimated variances of regression coefficients in misspecified error components models*. (Reprint Series). CentER for Economic Research.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

CBM
R

entER


8823

1992

for
Economic Research

NR. 76
76

REPRINT



On the Estimated Variances of
Regression Coefficients in
Misspecified Error Components
Models

by
Philippe J. Deschamps

518.931

Reprinted from *Econometric Theory*,
Vol. 7, No. 3, 1991



Reprint Series
no. 76

CENTER FOR ECONOMIC RESEARCH

Research Staff

Helmut Bester
Eric van Damme

Board

Helmut Bester
Eric van Damme, director
Arie Kapteyn

Scientific Council

Eduard Bomhoff	Erasmus University Rotterdam
Willem Buijer	Yale University
Jacques Drèze	Université Catholique de Louvain
Theo van de Klundert	Tilburg University
Simon Kuipers	Groningen University
Jean-Jacques Laffont	Université des Sciences Sociales de Toulouse
Merton Miller	University of Chicago
Stephen Nickell	University of Oxford
Pieter Ruys	Tilburg University
Jacques Sijben	Tilburg University

Residential Fellows

Svend Albaek	European University Institute
Pramila Krishnan	San Francisco State University
Jan Magnus	Tilburg University
Eduardo Siandra	UCLA
Dale Stahl II	University of Texas at Austin
Hideo Suehiro	Kobe University

Doctoral Students

Roel Beetsma
Hans Bloemen
Sjaak Hurkens
Frank de Jong
Pieter Kop Jansen

Address: Warandelaan 2, P.O. Box 90153, 5000 LE Tilburg, The Netherlands
Phone : +31 13 663050
Telex : 52426 kub nl
Telefax: +31 13 663066
E-mail : "center@htikub5.bitnet"

ISSN 0924-7874

1991

On the Estimated Variances of
Regression Coefficients in
Misspecified Error Components
Models

by
Philippe J. Deschamps

Reprinted from *Econometric Theory*,
Vol. 7, No. 3, 1991

Reprint Series
no. 76

ON THE ESTIMATED VARIANCES OF REGRESSION COEFFICIENTS IN MISSPECIFIED ERROR COMPONENTS MODELS

PHILIPPE J. DESCHAMPS
University of Fribourg

In a regression model with an arbitrary number of error components, the covariance matrix of the disturbances has three equivalent representations as linear combinations of matrices. Furthermore, this property is invariant with respect to powers, matrix addition, and matrix multiplication. This result is applied to the derivation and interpretation of the inconsistency of the estimated coefficient variances when the error components structure is improperly restricted. This inconsistency is defined as the difference between the asymptotic variance obtained when the restricted model is correctly specified, and the asymptotic variance obtained when the restricted model is incorrectly specified; when some error components are improperly omitted, and the remaining variance components are consistently estimated, it is always negative. In the case where the time component is improperly omitted from the two-way model, we show that the difference between the true and estimated coefficient variances is of order greater than N^{-1} in probability.

1. INTRODUCTION

Since the pioneering article by Balestra and Nerlove [3], many authors have studied the estimation of linear regression models with error components. The Balestra–Nerlove specification was limited to a two-way classification of data, and omitted time-specific effects. This restriction was later lifted by Wallace and Hussain [10] and Nerlove [8], who considered the following error specification:

$$u_{it} = v_i + w_t + \epsilon_{it} \quad \text{for } i = 1, \dots, n \quad \text{and} \quad t = 1, \dots, T \quad (1)$$

where v_i , w_t , and ϵ_{it} have zero expectations, constant variances, and are pairwise and serially uncorrelated. Nerlove, in particular, found the spectral form of the error covariance matrix V implied by equation (1), expressing V as a linear combination of symmetric, idempotent, and pairwise orthog-

Part of this research was conducted while the author held a Faculty Research Fellowship at CentER, Tilburg University. I wish to thank Pietro Balestra, Badi Baltagi, and anonymous referees for helpful comments. A previous version of this paper was presented at CORE, CentER, and at the European Meeting of the Econometric Society, Munich, 1989. Any remaining error is my own.

onal matrices that sum to an identity matrix. All the rational powers of V can be similarly represented; this is particularly useful for interpreting and computing the Aitken estimators of the regression coefficients. The spectral form also leads to a particularly simple expression of the ANOVA table for the variance components of u_{ii} (for a classical presentation of this table, see, e.g., Graybill [6, p. 349]).

More recently, Searle and Henderson [9] and Deschamps [4] have studied the structure of the error covariance matrix in a model with a general, p -way classification of data (where there are $2^p - 1$ error components). As will be shown, their results imply that V has three equivalent representations as linear combinations of matrices. The coefficients of the first one are the variance components; the coefficients of the second one (the spectral form) are the eigenvalues of V ; the coefficients of the third one are the elements of V . Each linear combination is, in a sense, isomorphic to the two others. Furthermore, the property is invariant with respect to powers, linear combinations, and matrix multiplication. This means that if the nonsingular matrices V_1, V_2, V_3, V_4 have the same form as V and if $a, b, \alpha, \beta, \gamma, \delta$ are scalars, then $V_1^\alpha (aV_2^\beta + bV_3^\gamma)V_4^\delta$ also has the same form; it can be represented as three linear combinations of the same matrices that appear in the three expressions for V . Furthermore, the three sets of coefficients are easily derived.

Section 2 of this paper will present the assumptions of the model, and a new notation. In Section 3, we present Theorem 1, which is a general result on the misspecified error components model. Theorem 1 can be used to derive and interpret the inconsistency ($\text{plim } N(\hat{\sigma}_i^2 - \sigma_i^2)$) of the estimated variances of the regression coefficients when some error components are improperly omitted (even though their variances are nonzero), and the remaining variance components are consistently estimated. We will show that the true coefficient variances are *always underestimated* in the misspecified model, for any p -way classification and for any number of omitted components. More generally, we find that the inconsistency can be expressed as three linear combinations of quadratic forms, in the same way as there are three representations of V . This turns out to be very helpful for the interpretation of the inconsistency.

When a *single* error component is omitted in the misspecified model and when the remaining variance components are consistently estimated, our results also imply that the inconsistency is *proportional* to the variance of the omitted component. The coefficient of proportionality is a linear combination of sums of squares of partial sums (as in, e.g., $\alpha_1 \sum_i (\sum_j Z_{ij})^2 + \alpha_2 \sum_i (\sum_j Z_{ij})^2$); it can be estimated as a by-product of generalized least squares on the misspecified model. This greatly facilitates the sensitivity analysis of misspecification.

Theorem 1 also applies to other types of misspecification: the inconsistency of an estimated coefficient variance is a linear combination of the inconsistencies of the variance component estimates.

In Section 4, we illustrate the previous results by analyzing the inconsistency when the time-specific component w_t is improperly omitted from equation (1). This case is of particular interest since it corresponds to the earliest (Balestra–Nerlove) specification, and is therefore found in most empirical implementations. In this instance the inconsistency has a particularly simple, intuitively appealing form, whose properties are easily investigated. It will be shown that this inconsistency is unbounded, unless the matrix of regressors satisfies very restrictive assumptions.

Lemma 1 of Appendix A presents the three isomorphic representations of V , and can be viewed as a synthesis of Searle and Henderson [9] and Deschamps [4]. This lemma will be used in Appendix B, where we prove Theorem 1 of Section 3. Appendix C presents estimators of the variance components which are consistent under misspecification, and coincide with the ones proposed by Amemiya [1] for a correctly specified three-component model.

2. A GENERAL ERROR COMPONENTS MODEL

Our regression model has 2^p error components, one of which is identically zero. It can be written as

$$y = X\beta + u \quad (2)$$

where y is an $N \times 1$ vector of observations on a dependent variable, X is an $N \times k$ matrix of observations on nonstochastic regressors, β is a $k \times 1$ vector of regression coefficients, and u is an $N \times 1$ vector of compound disturbances with the following structure:

$$u = \sum_{\alpha=00\dots 0}^{11\dots 1} (s_{n_1}^{1-\alpha_1} \otimes \dots \otimes s_{n_p}^{1-\alpha_p}) v_\alpha \quad (3a)$$

with

$$E(v_\alpha) = 0_{q_\alpha} \quad (3b)$$

$$E(v_\alpha v'_\alpha) = \theta_\alpha I_{q_\alpha} \quad (3c)$$

$$E(v_\alpha v'_\beta) = 0_{q_\alpha \times q_\beta} \quad (\alpha \neq \beta). \quad (3d)$$

In equations (3a)–(3d) and in the rest of this paper, the Greek *subscripts* α , β , and γ indicate *binary* numbers with p digits, for example, $\alpha = \alpha_1 \alpha_2 \dots \alpha_p$, where α_i is either zero or one. The letters n_1, \dots, n_p denote numbers of cells with $\prod_{i=1}^p n_i = N$, s_{n_i} is an $n_i \times 1$ vector of ones, and we adopt the convention that $s_{n_i}^0 = I_{n_i}$, an identity matrix of order n_i . v_α is the vector of the $q_\alpha = \prod_{i=1}^p n_i^{\alpha_i}$ realizations of error component α . This vector is assumed to have zero expectation and scalar covariance matrix in (3b) and (3c); equation (3d) assumes that the error components are pairwise uncorrelated. We take the *first* error component to be identically zero, and this implies that $\theta_{00\dots 0} = 0$.

Specification (3a)–(3d) obviously implies the following form for the covariance matrix of u (see Note 1):

$$V = E(uu') = \sum_{\alpha=00\dots 0}^{11\dots 1} \theta_{\alpha} (S_{n_1}^{1-\alpha_1} \otimes \dots \otimes S_{n_p}^{1-\alpha_p}) \quad (4)$$

where S_{n_i} is a square matrix of ones of order n_i and $S_{n_i}^0 \equiv I_{n_i}$.

The model in Nerlove [8] is obtained as a special case of (4) by letting $p = 2$, $n_1 = n$, and $n_2 = T$:

$$V = \theta_{00} S_{nT} + \theta_{01} (S_n \otimes I_T) + \theta_{10} (I_n \otimes S_T) + \theta_{11} I_{nT} \quad (5)$$

where $\theta_{00} = 0$, $\theta_{01} \equiv \sigma_w^2$, $\theta_{10} \equiv \sigma_v^2$, and $\theta_{11} \equiv \sigma_e^2$ (compare equation (1)). By letting $\theta_{01} = 0$ in (5), the model specializes further to Balestra and Nerlove [3].

The reader will have no difficulty in verifying that when $p = 3$, if $\theta_{000} = \theta_{001} = \theta_{010} = \theta_{011} = \theta_{101} = 0$ and if $\theta_{100} \equiv \sigma_v^2$, $\theta_{110} \equiv \sigma_w^2$, $\theta_{111} \equiv \sigma_e^2$, equation (4) specializes to the nested specification in Fuller and Battese [5], implied by

$$u_{ijt} = v_i + w_{ij} + \epsilon_{ijt}. \quad (6)$$

So the binary number α may serve to interpret the variance component θ_{α} in equation (4): it is simply the variance of that error term which has the particular index subset identified by the unitary digits in α .

Equation (4) is the first of our three isomorphic characterizations of V ; the two others are given in Appendix A.

3. THE CONSEQUENCES OF MISSPECIFICATION

We consider the case where V in equation (4) is estimated by:

$$\hat{V}_{\bullet} = \sum_{\alpha=00\dots 0}^{11\dots 1} \hat{\theta}_{\alpha}^* \left(\bigotimes_{i=1}^p S_{n_i}^{1-\alpha_i} \right) \quad (7)$$

where $\hat{\theta}^*$ is a possibly inconsistent estimator of the $2^p \times 1$ vector θ . We estimate β by

$$\hat{\beta} = (X' \hat{V}_{\bullet}^{-1} X)^{-1} X' \hat{V}_{\bullet}^{-1} y. \quad (8)$$

The covariance matrix of $\hat{\beta}$ is estimated by:

$$C = (X' \hat{V}_{\bullet}^{-1} X)^{-1}$$

whereas the true covariance matrix, conditional on $\hat{\theta}^*$, is

$$V(\hat{\beta} | \hat{\theta}^*) = (X' \hat{V}_{\bullet}^{-1} X)^{-1} X' \hat{V}_{\bullet}^{-1} V \hat{V}_{\bullet}^{-1} X (X' \hat{V}_{\bullet}^{-1} X)^{-1}.$$

We let $\theta^* = \text{plim } \hat{\theta}^*$ (see Note 2), and assume that $V_{\bullet} = \sum_{\alpha} \theta_{\alpha}^* \bigotimes_{i=1}^p S_{n_i}^{1-\alpha_i}$ is positive definite. We also assume that $X' V_{\bullet}^{-1} X / N$ tends to a finite, positive definite matrix M_{\bullet} . We further note that

$$NC = \left(\frac{X' \hat{V}_{\bullet}^{-1} X}{N} \right)^{-1}$$

and

$$NV(\hat{\beta}|\hat{\theta}^*) = \left(\frac{X' \hat{V}_*^{-1} X}{N} \right)^{-1} \left(\frac{X' \hat{V}_*^{-1} V \hat{V}_*^{-1} X}{N} \right) \left(\frac{X' \hat{V}_*^{-1} X}{N} \right)^{-1}$$

We wish to evaluate³

$$\begin{aligned} \text{plim } N(C - V(\hat{\beta}|\hat{\theta}^*)) \\ &= \text{plim } N((X' \hat{V}_*^{-1} X)^{-1} - (X' \hat{V}_*^{-1} X)^{-1} X' \hat{V}_*^{-1} V \hat{V}_*^{-1} X (X' \hat{V}_*^{-1} X)^{-1}) \\ &= \lim N((X' V_*^{-1} X)^{-1} - (X' V_*^{-1} X)^{-1} X' V_*^{-1} V V_*^{-1} X (X' V_*^{-1} X)^{-1}) \\ &= \lim N((X' V_*^{-1} X)^{-1} (X' V_*^{-1} V_* V_*^{-1} X - X' V_*^{-1} V V_*^{-1} X) (X' V_*^{-1} X)^{-1}) \\ &= M_*^{-1} \lim \left(\frac{X'_* D X_*}{N} \right) M_*^{-1} \end{aligned} \quad (9)$$

with $X_* = V_*^{-1/2} X$ and $D = V_*^{-1/2} (V_* - V) V_*^{-1/2}$.

Equation (9) implies the following expression for the inconsistency of the estimated variance of $\sqrt{N}\hat{\beta}_r$:

$$\text{plim } N(C_{rr} - V(\hat{\beta}_r|\hat{\theta}^*)) = \lim \left(\frac{Z'(\ell) D Z(\ell)}{N} \right) \quad (10)$$

where $Z(\ell)$ is the ℓ th column of the matrix $X_* M_*^{-1}$.

The following definition provides the key to the three equivalent representations of V given in Appendix A, and to the three corresponding representations of (10) given by Theorem 1 of this section.

DEFINITION 1. Let, as before, S_{n_i} be a matrix of ones of order n_i and zero powers denote identity matrices. We define

$$K = \bigotimes_{i=1}^p \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}; \quad (11)$$

$$K_0 = \bigotimes_{i=1}^p \begin{pmatrix} n_i & 1 \\ 0 & 1 \end{pmatrix}; \quad (12)$$

$$L_\beta = \bigotimes_{i=1}^p S_{n_i}^{1-\beta_i}; \quad (13)$$

$$R_\beta = \bigotimes_{i=1}^p \left(\frac{1}{n_i} S_{n_i} \right)^{1-\beta_i} \left(I_{n_i} - \frac{1}{n_i} S_{n_i} \right)^{\beta_i}; \quad (14)$$

$$M_\beta = \bigotimes_{i=1}^p (S_{n_i} - I_{n_i})^{1-\beta_i}. \quad (15) \quad \blacksquare$$

We now define $\eta = \theta^* - \theta$ as the vector of variance component inconsistencies. It is shown in Appendix A that the elements of $K_0 \theta^*$ are the eigenvalues of V_* , so that they are strictly positive by assumption. Theorem 1 shows that the inconsistency (10) of an estimated coefficient variance is a

linear combination of the elements of η , with coefficients that depend on θ^* , but not on θ .

THEOREM 1. *Let $\eta = \text{plim } \hat{\theta}^* - \theta = \theta^* - \theta$, let $V_* = \sum_{\alpha} \theta_{\alpha}^* L_{\alpha}$, let $D = V_*^{-1/2}(V_* - V)V_*^{-1/2}$, and let Δ be the diagonal matrix with the elements of $K_0\theta^*$ on the diagonal. Then D can be equivalently expressed as*

$$D = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_{\gamma} \sum_{\beta=00\dots 0}^{11\dots 1} a_{\beta\gamma} R_{\beta} \quad (16)$$

$$D = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_{\gamma} \sum_{\beta=00\dots 0}^{11\dots 1} b_{\beta\gamma} L_{\beta} \quad (17)$$

$$D = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_{\gamma} \sum_{\beta=00\dots 0}^{11\dots 1} c_{\beta\gamma} M_{\beta} \quad (18)$$

where $a_{\beta\gamma}$, $b_{\beta\gamma}$, and $c_{\beta\gamma}$ are the elements in row β and column γ of $\Delta^{-1}K_0$, $K_0^{-1}\Delta^{-1}K_0$, and $KK_0^{-1}\Delta^{-1}K_0$, respectively. ■

The most interesting type of misspecification occurs when one, or several, variance components θ_{γ} are improperly excluded from equation (4). This follows from letting $\hat{\theta}_{\gamma}^* = 0$, so that $\theta_{\gamma}^* = 0$. It is shown in Appendix C that the remaining variance components can always be consistently estimated. In this case the inconsistency (10) is always negative: indeed, it is shown in Appendix A that the matrices R_{β} in (16) are idempotent, pairwise orthogonal, and add up to an identity matrix, so that the eigenvalues of D are $\lambda_{\beta} = \sum_{\gamma} \eta_{\gamma} a_{\beta\gamma}$. Since $a_{\beta\gamma} \geq 0$ and since $\eta_{\gamma} = 0$ when component γ is included, whereas $\eta_{\gamma} = -\theta_{\gamma} \leq 0$ when component γ is excluded, we have that $\lambda_{\beta} \leq 0$ for all β .

We may also note that the opposite type of misspecification whereby a variance component θ_{γ} is improperly included is of no consequence asymptotically if $\text{plim } \hat{\theta}_{\gamma}^* = \theta_{\gamma} = 0$, so that $\eta_{\gamma} = 0$ in Theorem 1.

Equation (17) implies that the quadratic form in (10) is a linear combination of sums of squares of partial sums of the elements of $Z(\ell)$, as is obvious from the definition of L_{β} in (13). This fact will be used in Section 4. Similarly, equation (18) expresses the quadratic form as a linear combination of sums of cross-products of the elements of $Z(\ell)$ (see the definition of M_{β} in (15)). Using the arguments in Appendix A, it is easy to show that the coefficients $\sum_{\gamma} \eta_{\gamma} c_{\beta\gamma}$ in equation (18) are in fact the elements of D .

4. AN ILLUSTRATION

In this section, we illustrate the preceding theorem by considering the case where w_i is improperly omitted from equation (1); equivalently, $\hat{\theta}_{01}^* = 0$ in equation (7). We will first consider the case where the two remaining variance components are consistently estimated (Appendix C shows that this will be true if the two-way formulas are used). This assumption will be relaxed

at the end of this section. For now, we assume that $\theta_{00}^* = \theta_{01}^* = 0$, $\theta_{10}^* = \theta_{10}$, and $\theta_{11}^* = \theta_{11}$.

As previously noted, the elements of $K_0\theta^*$ in Theorem 1 are the eigenvalues of V_* . Using equation (12), they may be written as

$$\begin{pmatrix} \lambda_{00} \\ \lambda_{01} \\ \lambda_{10} \\ \lambda_{11} \end{pmatrix} = \begin{pmatrix} nT & n & T & 1 \\ 0 & n & 0 & 1 \\ 0 & 0 & T & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 0 \\ 0 \\ \theta_{10} \\ \theta_{11} \end{pmatrix} = \begin{pmatrix} T\theta_{10} + \theta_{11} \\ \theta_{11} \\ T\theta_{10} + \theta_{11} \\ \theta_{11} \end{pmatrix}. \quad (19)$$

Furthermore, the coefficients $b_{\beta\gamma}$ in equation (17) are elements of the matrix:

$$\begin{aligned} & K_0^{-1} \Delta^{-1} K_0 \\ &= \frac{1}{nT} \begin{pmatrix} 1 & -1 & -1 & 1 \\ 0 & T & 0 & -T \\ 0 & 0 & n & -n \\ 0 & 0 & 0 & nT \end{pmatrix} \begin{pmatrix} \lambda_{00}^{-1} & 0 & 0 & 0 \\ 0 & \lambda_{01}^{-1} & 0 & 0 \\ 0 & 0 & \lambda_{10}^{-1} & 0 \\ 0 & 0 & 0 & \lambda_{11}^{-1} \end{pmatrix} \begin{pmatrix} nT & n & T & 1 \\ 0 & n & 0 & 1 \\ 0 & 0 & T & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \\ &= \frac{1}{nT} \begin{pmatrix} nT\lambda_{00}^{-1} & n(\lambda_{00}^{-1} - \lambda_{01}^{-1}) & T(\lambda_{00}^{-1} - \lambda_{10}^{-1}) & \lambda_{00}^{-1} - \lambda_{01}^{-1} - \lambda_{10}^{-1} + \lambda_{11}^{-1} \\ 0 & nT\lambda_{01}^{-1} & 0 & T(\lambda_{01}^{-1} - \lambda_{11}^{-1}) \\ 0 & 0 & nT\lambda_{10}^{-1} & n(\lambda_{10}^{-1} - \lambda_{11}^{-1}) \\ 0 & 0 & 0 & nT\lambda_{11}^{-1} \end{pmatrix}. \end{aligned} \quad (20)$$

Since the only inconsistency occurs in the time component, we have $\eta_{01} = -\theta_{01}$ and $\eta_{00} = \eta_{10} = \eta_{11} = 0$, so that we only need those coefficients $b_{\beta\gamma}$ that are located in the second column of the preceding matrix (which has the binary index 01). Upon substituting (13), (19), and (20) in equation (17), we obtain

$$\begin{aligned} D &= -\theta_{01} \left(\frac{1}{T} \left(\frac{1}{T\theta_{10} + \theta_{11}} - \frac{1}{\theta_{11}} \right) S_{nT} + \frac{1}{\theta_{11}} (S_n \otimes I_T) \right) \\ &= -\frac{\theta_{01}}{\theta_{11}} \left(\frac{-\theta_{10}}{T\theta_{10} + \theta_{11}} S_{nT} + (S_n \otimes I_T) \right) \end{aligned} \quad (21)$$

so that the quadratic form in equation (10) is

$$Z'(\ell) D Z(\ell) = -\frac{\theta_{01}}{\theta_{11}} \left(\sum_{i=1}^T Z_{i\cdot}'(\ell) - \frac{\theta_{10}}{T\theta_{10} + \theta_{11}} Z_{\cdot\cdot}'(\ell) \right) \quad (22)$$

with $Z_{i\cdot}(\ell) = \sum_{\mu=1}^n Z_{i\mu}(\ell)$ and $Z_{\cdot\cdot}(\ell) = \sum_{i=1}^T \sum_{\mu=1}^n Z_{i\mu}(\ell)$.

Since $\theta_{10}/(T\theta_{10} + \theta_{11}) < 1/T$, an upper bound for the expression in (22) is given by

$$Z'(\ell)DZ(\ell) \leq -\frac{\theta_{01}}{\theta_{11}} \sum_{t=1}^T \left(Z_{.,t}(\ell) - \frac{Z_{..}(\ell)}{T} \right)^2$$

and it is seen that for given θ_{01} and θ_{11} , $\lim_{T \rightarrow \infty} Z'(\ell)DZ(\ell) = -\infty$, unless $\theta_{01} = 0$ or unless $Z_{.,t}(\ell) = \alpha$ for all t . In the latter case, equation (22) is easily shown to imply

$$Z'(\ell)DZ(\ell) = \frac{-T\alpha^2\theta_{01}}{T\theta_{10} + \theta_{11}}$$

whose limit equals $-\alpha^2\theta_{01}/\theta_{10}$ as T tends to infinity.

To summarize, the inconsistency of the ℓ th estimated variance in the misspecified model is unbounded as $n \rightarrow \infty$ and $T \rightarrow \infty$. This may be seen by dividing equation (22) by nT , and noting that $\sum_{t=1}^T Z_{.,t}^2(\ell)/nT$ does not converge in general. The unboundedness occurs unless the "observations" in the ℓ th column of $X_*M_*^{-1}$ repeat their average pattern over time. In this (very special) case where $Z_{.,t}(\ell) = \alpha$ for all t , we have

$$\text{plim } nT(C_{\ell\ell} - V(\hat{\beta}_\ell | \hat{\theta}^*)) = \lim \frac{Z'(\ell)DZ(\ell)}{nT} = \lim \frac{1}{nT} \left(\frac{-\alpha^2\theta_{01}}{\theta_{10}} \right) = 0.$$

We will now show that the previous result remains essentially unchanged when $\hat{\theta}_{10}^*$ and $\hat{\theta}_{11}^*$ are the one-way estimators:

$$\hat{\theta}_{11}^* = \frac{1}{n(T-1)} (\hat{u}'(I_n \otimes R_1)\hat{u})$$

$$\hat{\theta}_{10}^* = \frac{1}{T} \left(\frac{\hat{u}'(I_n \otimes R_0)\hat{u}}{n} - \frac{\hat{u}'(I_n \otimes R_1)\hat{u}}{n(T-1)} \right)$$

where $R_0 = T^{-1}S_T$, $R_1 = I_T - R_0$, and where \hat{u} is a predictor of u . If the true disturbances u_{it} are known and $\hat{u} = u$, it is easy to show that $E(\hat{\theta}_{11}^*) = \theta_{01} + \theta_{11}$ and $E(\hat{\theta}_{10}^*) = \theta_{10}$; we will therefore assume that $\theta_{11}^* = \theta_{01} + \theta_{11}$ and that $\theta_{10}^* = \theta_{10}$. As before, we have $\theta_{00}^* = \theta_{01}^* = 0$. The eigenvalues of V_* are again given by $K_0\theta^*$, this time as

$$\begin{pmatrix} \lambda_{00} \\ \lambda_{01} \\ \lambda_{10} \\ \lambda_{11} \end{pmatrix} = \begin{pmatrix} T\theta_{10} + \theta_{01} + \theta_{11} \\ \theta_{01} + \theta_{11} \\ T\theta_{10} + \theta_{01} + \theta_{11} \\ \theta_{01} + \theta_{11} \end{pmatrix}.$$

The vector η of variance component inconsistencies is this time given by:

$$\begin{pmatrix} \eta_{(0)} \\ \eta_{01} \\ \eta_{10} \\ \eta_{11} \end{pmatrix} = \begin{pmatrix} 0 \\ -\theta_{01} \\ 0 \\ \theta_{01} \end{pmatrix}.$$

Upon using again the methodology of Theorem 1, we see that $D = D_1 + D_2$, with

$$D_1 = -\theta_{01} \left(\frac{1}{T} \left(\frac{1}{T\theta_{10} + \theta_{01} + \theta_{11}} - \frac{1}{\theta_{01} + \theta_{11}} \right) S_{nT} + \frac{1}{\theta_{01} + \theta_{11}} (S_n \otimes I_T) \right)$$

$$D_2 = \theta_{01} \left(\frac{1}{T} \left(\frac{1}{T\theta_{10} + \theta_{01} + \theta_{11}} - \frac{1}{\theta_{01} + \theta_{11}} \right) (I_n \otimes S_T) + \frac{1}{\theta_{01} + \theta_{11}} I_{nT} \right).$$

The matrix D_1 has the same form as D in (21), with θ_{11} replaced by $\theta_{01} + \theta_{11}$; hence the preceding analysis also applies to $Z'(\ell)D_1Z(\ell)/nT$, which is negative and generally unbounded. By analogy with (22), the second term may be written as

$$\begin{aligned} Z'(\ell)D_2Z(\ell) &= \frac{\theta_{01}}{\theta_{01} + \theta_{11}} \left(\sum_{i=1}^n \sum_{t=1}^T Z_{it}^2(\ell) - \frac{\theta_{10}}{T\theta_{10} + \theta_{01} + \theta_{11}} \sum_{i=1}^n Z_{i.}^2(\ell) \right) \\ &\geq \frac{\theta_{01}}{\theta_{01} + \theta_{11}} \sum_{i=1}^n \sum_{t=1}^T \left(Z_{it}(\ell) - \frac{Z_{i.}(\ell)}{T} \right)^2 \geq 0 \end{aligned} \quad (23)$$

and $Z'(\ell)D_2Z(\ell)/nT$ converges to a finite, nonnegative number if $\sum_{i,t} Z_{it}^2(\ell)/nT$ and $\sum_i Z_{i.}^2(\ell)/nT^2$ are bounded. Furthermore the lower bound in (23) vanishes if, and only if, $Z_{it}(\ell) = c_i$ for all t ; in this case the limit of $Z'(\ell)D_2Z(\ell)/nT$ is easily shown to vanish when $\theta_{10} \neq 0$.

Collecting results, we may say that the inconsistency remains negative and generally unbounded, since the negative term dominates the positive one. Furthermore, when $\theta_{10} \neq 0$, the inconsistency now vanishes if (and only if) the elements of the relevant column of $X_*M_*^{-1}$ *exactly* repeat their pattern over time.

5. CONCLUDING REMARKS

In a general error components model with a p -way classification of data, we have given three interpretations of the inconsistency of the estimated coefficient variances when the error components structure is misspecified. In the special case where the time component is improperly omitted from the two-

way classification model, this inconsistency is in general unbounded, in the sense that $C_{\ell\ell} - V(\hat{\beta}_\ell | \hat{\theta}^*)$ is of order greater than N^{-1} in probability, so that $\text{plim } NV(\hat{\beta}_\ell | \hat{\theta}^*)$ does not exist. Since $\text{plim } NV(\hat{\beta}_\ell | \hat{\theta}^*)$ is the asymptotic variance when the restricted model is incorrectly specified, this implies that $\sqrt{N}(\hat{\beta}_\ell - \beta_\ell)$ does not have a proper limiting distribution in such a case.

The arguments in Appendixes A and C indicate that the p -way model is essentially a tensor generalization of the 2-way model, and is, as such, not much more difficult to handle. It is therefore unfortunate, in view of the potentially serious consequences of misspecification, that most contributions to the literature on error components have been limited to the Balestra-Nerlove specification.

NOTES

1. Note that equation (4) differs from Searle and Henderson [9, equation 2.2] in the indexing of the variance components θ_{α} . A correspondence between the two indexing schemes is obtained by writing the digits in α in reverse order, and taking the complement to unity of each digit: for instance, θ_{011} in our notation corresponds to θ_{001} in [9]. As we will see, our notation has definite advantages for interpreting the variance components, and for stating the isomorphisms of Appendix A.

2. Unless otherwise indicated, all the limits in this paper are taken as $n_i \rightarrow \infty$ for all i .

3. If $\hat{V}_\epsilon = \hat{\sigma}_\epsilon^2 I$ with $\hat{\sigma}_\epsilon^2$ the OLS estimator of the error variance, it may be shown (Greenwald, [7]) that $(X'X)^{-1}X'(\hat{V}_\epsilon - V)X(X'X)^{-1}$ approximates the bias of the estimated covariance matrix when N is large.

4. Note that $c_{\alpha\beta} = 0$ for $\beta = 00 \dots 0$.

REFERENCES

- Amemiya, T. The estimation of the variances in a variance components model. *International Economic Review* 12 (1971): 1-13.
- Balestra, P. Best quadratic unbiased estimators of the variance-covariance matrix in normal regression. *Journal of Econometrics* 1 (1973): 17-28.
- Balestra, P. & M. Nerlove. Pooling cross section and time series data in the estimation of a dynamic model: the demand for natural gas. *Econometrica* 34 (1966): 585-612.
- Deschamps, P. A note on isomorphic characterizations of the dispersion matrix in error components models. *Linear Algebra and Its Applications* 111 (1988): 147-150.
- Fuller, W.A. & G.E. Battese. Transformations for estimation of linear models with nested-error structure. *Journal of the American Statistical Association* 68 (1973): 626-632.
- Graybill, F.A. *An Introduction to Linear Statistical Models*, vol. 1. New York: McGraw-Hill, 1961.
- Greenwald, B.C. A general analysis of bias in the estimated standard errors of least squares coefficients. *Journal of Econometrics* 22 (1983): 323-338.
- Nerlove, M. A note on error components models. *Econometrica* 39 (1971): 383-396.
- Searle, S.R. & H.V. Henderson. Dispersion matrices for variance components models. *Journal of the American Statistical Association* 74 (1979): 465-470.
- Wallace, T.D. & A. Hussain. The use of error components models in combining cross section with time series data. *Econometrica* 37 (1969): 55-72.

APPENDIX A. CHARACTERIZATIONS OF THE ERROR COVARIANCE MATRIX

It is easily shown that the following lemma can be extended to all the rational powers of V , and also to linear combinations or products of matrices having the same structure as V (an example of such an extension is in fact given by Theorem 1).

LEMMA 1. *Let u have the specification (3a)–(3d), and let $V = E(uu')$. Then*

$$V = \sum_{\alpha=00\dots 0}^{11\dots 1} \theta_{\alpha} L_{\alpha} = \sum_{\alpha=00\dots 0}^{11\dots 1} \lambda_{\alpha} R_{\alpha} = \sum_{\alpha=00\dots 0}^{11\dots 1} \rho_{\alpha} M_{\alpha}$$

where L_{α} , R_{α} , and M_{α} are given by Definition 1. The λ_{α} are the eigenvalues of V and the ρ_{α} are the elements of V . Furthermore, if λ , ρ and θ denote the $2^p \times 1$ vectors with elements λ_{α} , ρ_{α} , and θ_{α} , we have $\lambda = K_0 \theta$ and $\rho = K \theta$, with K and K_0 given by (11) and (12).

Proof. Equation (4) may also be written as

$$V = (\theta' \otimes I_N) L \quad (24)$$

where

$$L = \begin{bmatrix} L_{00\dots 0} \\ \vdots \\ L_{11\dots 1} \end{bmatrix}$$

and where the matrices L_{α} are as defined in (13). Using $\lambda = K_0 \theta$ and $\rho = K \theta$, (24) is equivalent to

$$\begin{aligned} V &= (\theta' \otimes I_N) (K_0' \otimes I_N) ((K_0')^{-1} \otimes I_N) L \\ &= (\lambda' \otimes I_N) ((K_0')^{-1} \otimes I_N) L \equiv (\lambda' \otimes I_N) R \end{aligned} \quad (25)$$

and

$$\begin{aligned} V &= (\theta' \otimes I_N) (K' \otimes I_N) ((K')^{-1} \otimes I_N) L \\ &= (\rho' \otimes I_N) ((K')^{-1} \otimes I_N) L \equiv (\rho' \otimes I_N) M. \end{aligned} \quad (26)$$

If we partition R and M in the same fashion as L ,

$$R = \begin{bmatrix} R_{00\dots 0} \\ \vdots \\ R_{11\dots 1} \end{bmatrix} \quad \text{and} \quad M = \begin{bmatrix} M_{00\dots 0} \\ \vdots \\ M_{11\dots 1} \end{bmatrix},$$

it follows from (25) and (26) that

$$R_{\beta} = \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha}^0 L_{\alpha} \quad (27)$$

and

$$M_{\beta} = \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha} L_{\alpha} \quad (28)$$

where $k_{\beta\alpha}^0$ and $k_{\beta\alpha}$ are the elements in row β and column α of $(K'_0)^{-1}$ and $(K')^{-1}$, respectively.

We now show that (27) is the expansion of (14). When $p = 1$, we have

$$(K'_0)^{-1} = n_p^{-1} \begin{pmatrix} 1 & 0 \\ -1 & n_p \end{pmatrix}, \quad L_0 = S_{n_p}, \quad L_1 = I_{n_p},$$

so that (27) obviously implies (14). It is easy to see that the property is true for p whenever it is true for $p - 1$. Indeed, $(K'_0)^{-1} \equiv (K'_0(p))^{-1}$ and $L \equiv L(p)$ may be defined recursively as

$$(K'_0(p))^{-1} = \begin{pmatrix} n_1^{-1}(K'_0(p-1))^{-1} & O \\ -n_1^{-1}(K'_0(p-1))^{-1} & (K'_0(p-1))^{-1} \end{pmatrix}$$

$$L(p) = \begin{pmatrix} S_{n_1} \otimes L_{00\dots 0}(p-1) \\ \vdots \\ S_{n_1} \otimes L_{11\dots 1}(p-1) \\ I_{n_1} \otimes L_{00\dots 0}(p-1) \\ \vdots \\ I_{n_1} \otimes L_{11\dots 1}(p-1) \end{pmatrix}$$

so that, upon letting $\beta = (\beta_1 \gamma_1 \gamma_2 \dots \gamma_{p-1})$, (27) may be written if $\beta_1 = 0$ as

$$\begin{aligned} R_{\beta}(p) &= \sum_{\alpha} (n_1^{-1} k_{\gamma\alpha}^0(p-1))(S_{n_1} \otimes L_{\alpha}(p-1)) \\ &= n_1^{-1} S_{n_1} \otimes \sum_{\alpha} k_{\gamma\alpha}^0(p-1) L_{\alpha}(p-1) \\ &= n_1^{-1} S_{n_1} \otimes R_{\gamma}(p-1) \end{aligned}$$

and if $\beta_1 = 1$ as:

$$\begin{aligned} R_{\beta}(p) &= \sum_{\alpha} (-n_1^{-1} k_{\gamma\alpha}^0(p-1))(S_{n_1} \otimes L_{\alpha}(p-1)) \\ &\quad + \sum_{\alpha} (k_{\gamma\alpha}^0(p-1))(I_{n_1} \otimes L_{\alpha}(p-1)) \\ &= (I_{n_1} - n_1^{-1} S_{n_1}) \otimes \sum_{\alpha} k_{\gamma\alpha}^0(p-1) L_{\alpha}(p-1) \\ &= (I_{n_1} - n_1^{-1} S_{n_1}) \otimes R_{\gamma}(p-1) \end{aligned}$$

where we note that α and γ now involve $p - 1$ binary digits rather than p . The proof that (28) is the expansion of (15) is exactly similar.

We must show that the matrices R_{β} in (14) are idempotent, pairwise orthogonal, and add up to an identity matrix, thus proving our claim that λ is indeed the vector

of eigenvalues. $R_\beta R_\beta = R_\beta$ is obvious since R_β is a Kronecker product of idempotent matrices. $R_\alpha R_\beta = O$ for $\alpha \neq \beta$ follows from the fact that $n_i^{-1} S_{n_i} (I_{n_i} - n_i^{-1} S_{n_i}) = O$; indeed, $\alpha \neq \beta$ implies the existence of an index i such that $\alpha_i = 0$ and $\beta_i = 1$, or $\alpha_i = 1$ and $\beta_i = 0$. So $\sum_\beta R_\beta$ is symmetric and idempotent; in order to show that it equals I_N , we show that its trace is nonzero. This follows easily from (27), which implies

$$\text{tr} \left(\sum_{\beta=00\dots 0}^{11\dots 1} R_\beta \right) = \sum_{\beta=00\dots 0}^{11\dots 1} \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha}^0 \text{tr}(L_\alpha) = N \sum_{\beta=00\dots 0}^{11\dots 1} \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha}^0 = N$$

since $\sum_\beta \sum_\alpha k_{\beta\alpha}^0$ is the sum of the elements of $(K'_0)^{-1}$, which is easily seen to equal unity:

$$s'(K'_0)^{-1}s = \left(\bigotimes_{i=1}^p (1 \quad 1) \right) \left(\bigotimes_{i=1}^p \begin{pmatrix} n_i^{-1} & 0 \\ -n_i^{-1} & 1 \end{pmatrix} \right) \left(\bigotimes_{i=1}^p \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) = 1.$$

It was shown in Deschamps [4] that the elements of ρ are also the elements of V . Since, from (15), the elements of M_β are 0 or 1, this is also implied by $\sum_\beta M_\beta = S_N$, which is readily verified from (28).

APPENDIX B. PROOF OF THEOREM 1

It is an immediate consequence of Lemma 1 that V , V_* , and $V_* - V$ can be expressed as (different) linear combinations of the same matrices R_α . The coefficients in the linear combinations are the eigenvalues; in the case of $V_* - V$, they form the vector $K_0\theta^* - K_0\theta = K_0\eta$.

As is easily seen from the properties of the matrices R_α , both $V_*^{-1/2}$ and D can again be represented as linear combinations of the same matrices R_α . The eigenvalues of D are obtained by multiplying the eigenvalues of $V_* - V$ by those of V_*^{-1} , so that they form the vector $\Delta^{-1}K_0\eta$, with elements $\sum_\gamma \eta_\gamma a_{\beta\gamma}$. This proves that (16) is indeed the spectral form of D .

In order to obtain equation (17), we substitute (27) into (16):

$$D = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_\gamma \sum_{\beta=00\dots 0}^{11\dots 1} a_{\beta\gamma} \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha}^0 L_\alpha = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_\gamma \sum_{\alpha=00\dots 0}^{11\dots 1} \left(\sum_{\beta=00\dots 0}^{11\dots 1} k_{\beta\alpha}^0 a_{\beta\gamma} \right) L_\alpha$$

which is equivalent to (17) since $k_{\beta\alpha}^0$ is the element in row α and column β of K_0^{-1} .

In order to obtain equation (18), we note that (25) and (26) imply $R = ((K'_0)^{-1} \otimes I_N)L$ and $L = (K' \otimes I_N)M$, so that

$$R = ((K'_0)^{-1} K' \otimes I_N)M$$

and we may write, similarly to (27) and (28),

$$R_\beta = \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha}^* M_\alpha \quad (29)$$

where $k_{\beta\alpha}^*$ is the element in row β and column α of $(KK_0^{-1})'$. Upon substituting (29) into (16), we obtain

$$D = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_{\gamma} \sum_{\beta=00\dots 0}^{11\dots 1} a_{\beta\gamma} \sum_{\alpha=00\dots 0}^{11\dots 1} k_{\beta\alpha}^* M_{\alpha} = \sum_{\gamma=00\dots 0}^{11\dots 1} \eta_{\gamma} \sum_{\alpha=00\dots 0}^{11\dots 1} \left(\sum_{\beta=00\dots 0}^{11\dots 1} k_{\beta\alpha}^* a_{\beta\gamma} \right) M_{\alpha}$$

and (18) follows upon noting that $k_{\beta\alpha}^*$ is the element in row α and column β of KK_0^{-1} .

APPENDIX C. ESTIMATION OF THE VARIANCE COMPONENTS

This appendix may be viewed as a generalization of the first part of Amemiya [1] to the p -way classification; however, the method of proof is partly new and relies on Lemma 1. We first note that, since $\theta = K_0^{-1}\lambda$, unbiased estimators of the variance components are easy to obtain from unbiased estimators of the eigenvalues; however, since $\theta_{00\dots 0} = 0$ is known and since K_0^{-1} is upper triangular, we need only estimate the last $2^p - 1$ eigenvalues. For this reason we will always assume in this appendix that $\alpha \neq 00\dots 0$.

We will first present unbiased estimators $\tilde{\lambda}_{\alpha}$ of the eigenvalues when the true disturbances are known. The $2^p - 1$ last equations of $\tilde{\theta} = K_0^{-1}\tilde{\lambda}$ then provide unbiased estimators of the variance components θ_{α} .

We will show that $\tilde{\theta}_{\alpha}$ is consistent under the normality of u by proving that $V(\tilde{\theta}_{\alpha}) \rightarrow 0$; the usual continuity argument does not apply since K_0^{-1} is a function of the numbers of observations n_i . We will then show that predicting the true disturbances by the analysis of covariance residuals (a procedure suggested by Amemiya) leads to estimated variance components $\hat{\theta}_{\alpha}$ with the same asymptotic distribution as the $\tilde{\theta}_{\alpha}$. It follows that $\hat{\theta}_{\alpha}$ is a consistent, and asymptotically unbiased, estimator of θ_{α} .

In the omitted component case of Section 3, we let $\hat{\theta}_{\gamma}^* = 0$, and $\hat{\theta}_{\alpha}^* = \hat{\theta}_{\alpha}$ for $\alpha \neq \gamma$; misspecification does not affect the consistency of $\hat{\theta}_{\alpha}^*$ since λ is estimated from the correctly specified model.

We first show that when the true disturbances are known, unbiased estimators of the eigenvalues are given by $\tilde{\lambda}_{\alpha} = \mu_{\alpha}^{-1} u' R_{\alpha} u$, where, from (14), $\mu_{\alpha} = \text{tr}(R_{\alpha}) = \prod_{i=1}^p (n_i - 1)^{n_i}$. This follows from Balestra [2, Lemma 1]:

$$E(\tilde{\lambda}_{\alpha}) = \mu_{\alpha}^{-1} \text{tr}(R_{\alpha} V) = \mu_{\alpha}^{-1} \text{tr} \left(R_{\alpha} \sum_{\beta} \lambda_{\beta} R_{\beta} \right) = \lambda_{\alpha} \mu_{\alpha}^{-1} \text{tr}(R_{\alpha}) = \lambda_{\alpha}.$$

Under the normality of u , we have, from Balestra [2, Lemma 6]:

$$V(u' R_{\alpha} u) = 2 \text{tr}(R_{\alpha} V R_{\alpha} V) = 2 \text{tr}(\lambda_{\alpha}^2 R_{\alpha}) = 2 \lambda_{\alpha}^2 \mu_{\alpha};$$

furthermore $u' R_{\alpha} u$ and $u' R_{\beta} u$ are independent for $\alpha \neq \beta$. Upon letting $\tilde{\theta}_{\alpha} = \sum_{\beta} c_{\alpha\beta} \tilde{\lambda}_{\beta}$ with $c_{\alpha\beta}$ the element in row α and column β of K_0^{-1} , we then have

$$V(\tilde{\theta}_{\alpha}) = 2 \sum_{\beta} c_{\alpha\beta}^2 \mu_{\beta}^{-1} \lambda_{\beta}^2 = 2 \lambda' K_0' \Delta \Delta_{\alpha} K_0 \theta \quad (30)$$

where Δ and Δ_α are diagonal matrices with the μ_β^{-1} and the $c_{\alpha\beta}^2$ on the diagonal, respectively. It may be checked that

$$\Delta = \frac{1}{\prod_{i=1}^p (n_i - 1)} \bigotimes_{i=1}^p \begin{pmatrix} n_i - 1 & 0 \\ 0 & 1 \end{pmatrix} \quad (31)$$

$$\Delta_\alpha = \frac{1}{N^2} \bigotimes_{i=1}^p \begin{pmatrix} 0 & 0 \\ 0 & n_i^2 \end{pmatrix}^{\alpha_i} \quad (32)$$

so that the matrix in the quadratic form (30) may be written

$$\begin{aligned} K'_0 \Delta \Delta_\alpha K_0 &= \frac{1}{N^2 \prod_{i=1}^p (n_i - 1)} \bigotimes_{i=1}^p \begin{pmatrix} n_i & 0 \\ 1 & 1 \end{pmatrix} \bigotimes_{i=1}^p \begin{pmatrix} n_i - 1 & 0 \\ 0 & 1 \end{pmatrix} \\ &\quad \bigotimes_{i=1}^p \begin{pmatrix} 0 & 0 \\ 0 & n_i^2 \end{pmatrix}^{\alpha_i} \bigotimes_{i=1}^p \begin{pmatrix} n_i & 1 \\ 0 & 1 \end{pmatrix} \\ &= \frac{1}{N^2 \prod_{i=1}^p (n_i - 1)} \bigotimes_{i=1}^p \begin{pmatrix} 0 & 0 \\ 0 & n_i^2 \end{pmatrix}^{\alpha_i} \begin{pmatrix} n_i^2(n_i - 1) & n_i(n_i - 1)^{1-\alpha_i} \\ n_i(n_i - 1) & n_i \end{pmatrix} \\ &= \bigotimes_{i=1}^p \begin{pmatrix} 0 & 0 \\ 0 & (n_i - 1)^{-1} \end{pmatrix}^{\alpha_i} \begin{pmatrix} 1 & n_i^{-1} \\ n_i^{-1} & n_i^{-1}(n_i - 1)^{-1} \end{pmatrix}^{1-\alpha_i} \stackrel{\text{def}}{=} P_\alpha. \end{aligned} \quad (33)$$

It is easily seen from (33) that $\lim P_\alpha = O$, so that $V(\tilde{\theta}_\alpha) \rightarrow 0$ in (30). Furthermore, it can be checked that $\lim \mu_\alpha P_\alpha = e_\alpha e'_\alpha$, where e_α is the $2^p \times 1$ unit vector with element α equal to unity; equation (30) then implies $\lim \mu_\alpha V(\tilde{\theta}_\alpha) = 2\theta_\alpha^2$. This result, as well as equations (30) and (33), is consistent with Amemiya [1, pp. 5 and 6].

We now show the asymptotic equivalence when u is replaced by $\hat{u} = My = Mu$, where

$$M = I_N - \frac{1}{N} S_N - \left(I_N - \frac{1}{N} S_N \right) (X(X'QX)^{-1}X'Q)$$

with $Q \equiv R_{11 \dots 1} = \bigotimes_{i=1}^p (I_{n_i} - n_i^{-1} S_{n_i})$ (see Amemiya [1]). Since $R_\alpha S_N = S_N R_\alpha = O$, we have

$$M'R_\alpha M = R_\alpha - Q'X(X'QX)^{-1}X'R_\alpha X(X'QX)^{-1}X'Q.$$

Hence, upon letting

$$\hat{\theta}_\alpha = \sum_\beta c_{\alpha\beta} \hat{\lambda}_\beta = \sum_\beta c_{\alpha\beta} \mu_\beta^{-1} \hat{u}' R_\beta \hat{u}$$

we see that⁴

$$\begin{aligned} \sqrt{\mu_\alpha}(\hat{\theta}_\alpha - \theta_\alpha) &= \sqrt{\mu_\alpha}(\tilde{\theta}_\alpha - \theta_\alpha) \\ &\quad - \sqrt{\mu_\alpha} \sum_\beta c_{\alpha\beta} \mu_\beta^{-1} \frac{u' Q X}{\sqrt{N}} \left(\frac{X' Q X}{N} \right)^{-1} \left(\frac{X' R_\beta X}{N} \right) \left(\frac{X' Q X}{N} \right)^{-1} \frac{X' Q u}{\sqrt{N}}. \end{aligned} \quad (34)$$

The estimators $\hat{\theta}_\alpha$ and $\bar{\theta}_\alpha$ will be asymptotically equivalent if the last terms in (34) converge in probability to zero. If $(X'QX)/N$ tends to a finite, positive definite matrix and if $(X'R_\beta X)/N$ is bounded, this will be true when $\sqrt{\mu_\alpha} c_{\alpha\beta} \mu_\beta^{-1}$ tends to 0 for all β , by the same argument as in Amemiya [1, equation (16)]. But $c_{\alpha\beta} \mu_\beta^{-1}$ is the element in row α and column β of $K_0^{-1}\Delta$, where Δ is defined in (31). Since, from (12) and (31),

$$K_0^{-1}\Delta = \frac{1}{N \prod_{i=1}^p (n_i - 1)} \bigotimes_{i=1}^p \begin{pmatrix} 1 & -1 \\ 0 & n_i \end{pmatrix} \begin{pmatrix} n_i - 1 & 0 \\ 0 & 1 \end{pmatrix} = \frac{1}{N} \bigotimes_{i=1}^p \begin{pmatrix} 1 & -(n_i - 1)^{-1} \\ 0 & n_i (n_i - 1)^{-1} \end{pmatrix},$$

we see that this is indeed the case.

- No. 1 G. Marini and F. van der Ploeg, Monetary and fiscal policy in an optimising model with capital accumulation and finite lives, The Economic Journal, vol. 98, no. 392, 1988, pp. 772 - 786.
- No. 2 F. van der Ploeg, International policy coordination in interdependent monetary economies, Journal of International Economics, vol. 25, 1988, pp. 1 - 23.
- No. 3 A.P. Barten, The history of Dutch macroeconomic modelling (1936-1986), in W. Driehuis, M.M.G. Fase and H. den Hartog (eds.), Challenges for Macroeconomic Modelling, Contributions to Economic Analysis 178, Amsterdam: North-Holland, 1988, pp. 39 - 88.
- No. 4 F. van der Ploeg, Disposable income, unemployment, inflation and state spending in a dynamic political-economic model, Public Choice, vol. 60, 1989, pp. 211 - 239.
- No. 5 Th. ten Raa and F. van der Ploeg, A statistical approach to the problem of negatives in input-output analysis, Economic Modelling, vol. 6, no. 1, 1989, pp. 2 - 19.
- No. 6 E. van Damme, Renegotiation-proof equilibria in repeated prisoners' dilemma, Journal of Economic Theory, vol. 47, no. 1, 1989, pp. 206 - 217.
- No. 7 C. Mulder and F. van der Ploeg, Trade unions, investment and employment in a small open economy: a Dutch perspective, in J. Muysken and C. de Neubourg (eds.), Unemployment in Europe, London: The MacMillan Press Ltd, 1989, pp. 200 - 229.
- No. 8 Th. van de Klundert and F. van der Ploeg, Wage rigidity and capital mobility in an optimizing model of a small open economy, De Economist 137, nr. 1, 1989, pp. 47 - 75.
- No. 9 G. Dhaene and A.P. Barten, When it all began: the 1936 Tinbergen model revisited, Economic Modelling, vol. 6, no. 2, 1989, pp. 203 - 219.
- No. 10 F. van der Ploeg and A.J. de Zeeuw, Conflict over arms accumulation in market and command economies, in F. van der Ploeg and A.J. de Zeeuw (eds.), Dynamic Policy Games in Economics, Contributions to Economic Analysis 181, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1989, pp. 91 - 119.
- No. 11 J. Driffill, Macroeconomic policy games with incomplete information: some extensions, in F. van der Ploeg and A.J. de Zeeuw (eds.), Dynamic Policy Games in Economics, Contributions to Economic Analysis 181, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1989, pp. 289 - 322.
- No. 12 F. van der Ploeg, Towards monetary integration in Europe, in P. De Grauwe e.a., De Europese Monetair Integratie: vier visies, Wetenschappelijke Raad voor het Regeringsbeleid V 66, 's-Gravenhage: SDU uitgeverij, 1989, pp. 81 - 106.

- No. 13 R.J.M. Alessie and A. Kapteyn, Consumption, savings and demography, in A. Wenig, K.F. Zimmermann (eds.), Demographic Change and Economic Development, Berlin/Heidelberg: Springer-Verlag, 1989, pp. 272 - 305.
- No. 14 A. Hoque, J.R. Magnus and B. Pesaran, The exact multi-period mean-square forecast error for the first-order autoregressive model, Journal of Econometrics, vol. 39, no. 3, 1988, pp. 327 - 346.
- No. 15 R. Alessie, A. Kapteyn and B. Melenberg, The effects of liquidity constraints on consumption: estimation from household panel data, European Economic Review 33, no. 2/3, 1989, pp. 547 - 555.
- No. 16 A. Holly and J.R. Magnus, A note on instrumental variables and maximum likelihood estimation procedures, Annales d'Économie et de Statistique, no. 10, April-June, 1988, pp. 121 - 138.
- No. 17 P. ten Hacken, A. Kapteyn and I. Woittiez, Unemployment benefits and the labor market, a micro/macro approach, in B.A. Gustafsson and N. Anders Klevmarken (eds.), The Political Economy of Social Security, Contributions to Economic Analysis 179, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1989, pp. 143 - 164.
- No. 18 T. Wansbeek and A. Kapteyn, Estimation of the error-components model with incomplete panels, Journal of Econometrics, vol. 41, no. 3, 1989, pp. 341 - 361.
- No. 19 A. Kapteyn, P. Kooreman and R. Willemsse, Some methodological issues in the implementation of subjective poverty definitions, The Journal of Human Resources, vol. 23, no. 2, 1988, pp. 222 - 242.
- No. 20 Th. van de Klundert and F. van der Ploeg, Fiscal policy and finite lives in interdependent economies with real and nominal wage rigidity, Oxford Economic Papers, vol. 41, no. 3, 1989, pp. 459 - 489.
- No. 21 J.R. Magnus and B. Pesaran, The exact multi-period mean-square forecast error for the first-order autoregressive model with an intercept, Journal of Econometrics, vol. 42, no. 2, 1989, pp. 157 - 179.
- No. 22 F. van der Ploeg, Two essays on political economy: (i) The political economy of overvaluation, The Economic Journal, vol. 99, no. 397, 1989, pp. 850 - 855; (ii) Election outcomes and the stockmarket, European Journal of Political Economy, vol. 5, no. 1, 1989, pp. 21 - 30.
- No. 23 J.R. Magnus and A.D. Woodland, On the maximum likelihood estimation of multivariate regression models containing serially correlated error components, International Economic Review, vol. 29, no. 4, 1988, pp. 707 - 725.
- No. 24 A.J.J. Talman and Y. Yamamoto, A simplicial algorithm for stationary point problems on polytopes, Mathematics of Operations Research, vol. 14, no. 3, 1989, pp. 383 - 399.

- No. 25 E. van Damme, Stable equilibria and forward induction, Journal of Economic Theory, vol. 48, no. 2, 1989, pp. 476 - 496.
- No. 26 A.P. Barten and L.J. Bettendorf, Price formation of fish: An application of an inverse demand system, European Economic Review, vol. 33, no. 8, 1989, pp. 1509 - 1525.
- No. 27 G. Noldeke and E. van Damme, Signalling in a dynamic labour market, Review of Economic Studies, vol. 57 (1), no. 189, 1990, pp. 1 - 23
- No. 28 P. Kop Jansen and Th. ten Raa, The choice of model in the construction of input-output coefficients matrices, International Economic Review, vol. 31, no. 1, 1990, pp. 213 - 227.
- No. 29 F. van der Ploeg and A.J. de Zeeuw, Perfect equilibrium in a model of competitive arms accumulation, International Economic Review, vol. 31, no. 1, 1990, pp. 131 - 146.
- No. 30 J.R. Magnus and A.D. Woodland, Separability and aggregation, Economica, vol. 57, no. 226, 1990, pp. 239 - 247.
- No. 31 F. van der Ploeg, International interdependence and policy coordination in economies with real and nominal wage rigidity, Greek Economic Review, vol. 10, no. 1, June 1988, pp. 1 - 48.
- No. 32 E. van Damme, Signaling and forward induction in a market entry context, Operations Research Proceedings 1989, Berlin-Heidelberg: Springer-Verlag, 1990, pp. 45 - 59.
- No. 33 A.P. Barten, Toward a levels version of the Rotterdam and related demand systems, Contributions to Operations Research and Economics, Cambridge: MIT Press, 1989, pp. 441 - 465.
- No. 34 F. van der Ploeg, International coordination of monetary policies under alternative exchange-rate regimes, Advanced Lectures in Quantitative Economics, London-Orlando: Academic Press Ltd., 1990, pp. 91 - 121.
- No. 35 Th. van de Klundert, On socioeconomic causes of 'wait unemployment', European Economic Review, vol. 34, no. 5, 1990, pp. 1011 - 1022.
- No. 36 R.J.M. Alessie, A. Kapteyn, J.B. van Lochem and T.J. Wansbeek, Individual effects in utility consistent models of demand, in J. Hartog, G. Ridder and J. Theeuwes (eds.), Panel Data and Labor Market Studies, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1990, pp. 253 - 278.
- No. 37 F. van der Ploeg, Capital accumulation, inflation and long-run conflict in international objectives, Oxford Economic Papers, vol. 42, no. 3, 1990, pp. 501 - 525.
- No. 38 Th. Nijman and F. Palm, Parameter identification in ARMA Processes in the presence of regular but incomplete sampling, Journal of Time Series Analysis, vol. 11, no. 3, 1990, pp. 239 - 248.

- No. 39 Th. van de Klundert, Wage differentials and employment in a two-sector model with a dual labour market, Metroeconomica, vol. 40, no. 3, 1989, pp. 235 - 256.
- No. 40 Th. Nijman and M.F.J. Steel, Exclusion restrictions in instrumental variables equations, Econometric Reviews, vol. 9, no. 1, 1990, pp. 37 - 55.
- No. 41 A. van Soest, I. Woittiez and A. Kapteyn, Labor supply, income taxes, and hours restrictions in the Netherlands, Journal of Human Resources, vol. 25, no. 3, 1990, pp. 517 - 558.
- No. 42 Th.C.M.J. van de Klundert and A.B.T.M. van Schaik, Unemployment persistence and loss of productive capacity: a Keynesian approach, Journal of Macroeconomics, vol. 12, no. 3, 1990, pp. 363 - 380.
- No. 43 Th. Nijman and M. Verbeek, Estimation of time-dependent parameters in linear models using cross-sections, panels, or both, Journal of Econometrics, vol. 46, no. 3, 1990, pp. 333 - 346.
- No. 44 E. van Damme, R. Selten and E. Winter, Alternating bid bargaining with a smallest money unit, Games and Economic Behavior, vol. 2, no. 2, 1990, pp. 188 - 201.
- No. 45 C. Dang, The D_n -triangulation of \mathbb{T}^n for simplicial algorithms for computing solutions of nonlinear equations, Mathematics of Operations Research, vol. 16, no. 1, 1991, pp. 148 - 161.
- No. 46 Th. Nijman and F. Palm, Predictive accuracy gain from disaggregate sampling in ARIMA models, Journal of Business & Economic Statistics, vol. 8, no. 4, 1990, pp. 405 - 415.
- No. 47 J.R. Magnus, On certain moments relating to ratios of quadratic forms in normal variables: further results, Sankhya: The Indian Journal of Statistics, vol. 52, series B, part. 1, 1990, pp. 1 - 13.
- No. 48 M.F.J. Steel, A Bayesian analysis of simultaneous equation models by combining recursive analytical and numerical approaches, Journal of Econometrics, vol. 48, no. 1/2, 1991, pp. 83 - 117.
- No. 49 F. van der Ploeg and C. Withagen, Pollution control and the ramsey problem, Environmental and Resource Economics, vol. 1, no. 2, 1991, pp. 215 - 236.
- No. 50 F. van der Ploeg, Money and capital in interdependent economies with overlapping generations, Economica, vol. 58, no. 230, 1991, pp. 233 - 256.
- No. 51 A. Kapteyn and A. de Zeeuw, Changing incentives for economic research in the Netherlands, European Economic Review, vol. 35, no. 2/3, 1991, pp. 603 - 611.
- No. 52 C.G. de Vries, On the relation between GARCH and stable processes, Journal of Econometrics, vol. 48, no. 3, 1991, pp. 313 - 324.

- No. 53 R. Alessie and A. Kapteyn, Habit formation, interdependent preferences and demographic effects in the almost ideal demand system, The Economic Journal, vol. 101, no. 406, 1991, pp. 404 - 419.
- No. 54 W. van Groenendaal and A. de Zeeuw, Control, coordination and conflict on international commodity markets, Economic Modelling, vol. 8, no. 1, 1991, pp. 90 - 101.
- No. 55 F. van der Ploeg and A.J. Markink, Dynamic policy in linear models with rational expectations of future events: A computer package, Computer Science in Economics and Management, vol. 4, no. 3, 1991, pp. 175 - 199.
- No. 56 H.A. Keuzenkamp and F. van der Ploeg, Savings, investment, government finance, and the current account: The Dutch experience, in G. Alogoskoufis, L. Papademos and R. Portes (eds.), External Constraints on Macroeconomic Policy: The European Experience, Cambridge: Cambridge University Press, 1991, pp. 219 - 263.
- No. 57 Th. Nijman, M. Verbeek and A. van Soest, The efficiency of rotating-panel designs in an analysis-of-variance model, Journal of Econometrics, vol. 49, no. 3, 1991, pp. 373 - 399.
- No. 58 M.F.J. Steel and J.-F. Richard, Bayesian multivariate exogeneity analysis - an application to a UK money demand equation, Journal of Econometrics, vol. 49, no. 1/2, 1991, pp. 239 - 274.
- No. 59 Th. Nijman and F. Palm, Generalized least squares estimation of linear models containing rational future expectations, International Economic Review, vol. 32, no. 2, 1991, pp. 383 - 389.
- No. 60 E. van Damme, Equilibrium selection in 2×2 games, Revista Espanola de Economia, vol. 8, no. 1, 1991, pp. 37 - 52.
- No. 61 E. Bennett and E. van Damme, Demand commitment bargaining: the case of apex games, in R. Selten (ed.), Game Equilibrium Models III - Strategic Bargaining, Berlin: Springer-Verlag, 1991, pp. 118 - 140.
- No. 62 W. Güth and E. van Damme, Gorbby games - a game theoretic analysis of disarmament campaigns and the defense efficiency - hypothesis -, in R. Avenhaus, H. Karkar and M. Rudnianski (eds.), Defense Decision Making - Analytical Support and Crisis Management, Berlin: Springer-Verlag, 1991, pp. 215 - 240.
- No. 63 A. Roell, Dual-capacity trading and the quality of the market, Journal of Financial Intermediation, vol. 1, no. 2, 1990, pp. 105 - 124.
- No. 64 Y. Dai, G. van der Laan, A.J.J. Talman and Y. Yamamoto, A simplicial algorithm for the nonlinear stationary point problem on an unbounded polyhedron, Siam Journal of Optimization, vol. 1, no. 2, 1991, pp. 151 - 165.
- No. 65 M. McAleer and C.R. McKenzie, Keynesian and new classical models of unemployment revisited, The Economic Journal, vol. 101, no. 406, 1991, pp. 359 - 381.

- No. 66 A.J.J. Talman, General equilibrium programming, Nieuw Archief voor Wiskunde, vol. 8, no. 3, 1990, pp. 387 - 397.
- No. 67 J.R. Magnus and B. Pesaran, The bias of forecasts from a first-order autoregression, Econometric Theory, vol. 7, no. 2, 1991, pp. 222 - 235.
- No. 68 F. van der Ploeg, Macroeconomic policy coordination issues during the various phases of economic and monetary integration in Europe, European Economy - The Economics of EMU, Commission of the European Communities, special edition no. 1, 1991, pp. 136 - 164.
- No. 69 H. Keuzenkamp, A precursor to Muth: Tinbergen's 1932 model of rational expectations, The Economic Journal, vol. 101, no. 408, 1991, pp. 1245 - 1253.
- No. 70 L. Zou, The target-incentive system vs. the price-incentive system under adverse selection and the ratchet effect, Journal of Public Economics, vol. 46, no. 1, 1991, pp. 51 - 89.
- No. 71 E. Bomhoff, Between price reform and privatization: Eastern Europe in transition, Finanzmarkt und Portfolio Management, vol. 5, no. 3, 1991, pp. 241 - 251.
- No. 72 E. Bomhoff, Stability of velocity in the major industrial countries: a Kalman filter approach, International Monetary Fund Staff Papers, vol. 38, no. 3, 1991, pp. 626 - 642.
- No. 73 E. Bomhoff, Currency convertibility: when and how? A contribution to the Bulgarian debate, Kredit und Kapital, vol. 24, no. 3, 1991, pp. 412 - 431.
- No. 74 H. Keuzenkamp and F. van der Ploeg, Perceived constraints for Dutch unemployment policy, in C. de Neubourg (ed.), The Art of Full Employment - Unemployment Policy in Open Economies, Contributions to Economic Analysis 203, Amsterdam: Elsevier Science Publishers B.V. (North-Holland), 1991, pp. 7 - 37.
- No. 75 H. Peters and E. van Damme, Characterizing the Nash and Raiffa bargaining solutions by disagreement point axioms, Mathematics of Operations Research, vol. 16, no. 3, 1991, pp. 447 - 461.
- No. 76 P.J. Deschamps, On the estimated variances of regression coefficients in misspecified error components models, Econometric Theory, vol. 7, no. 3, 1991, pp. 369 - 384.

P.O. BOX 90153 5000 LE TILBURG THE NETHERLANDS

Bibliotheek K. U. Brabant



17 000 01169739 9